



# **SNOMED CT in Pathology**

**Marcial García Rojo**

Hospital General Universitario de Ciudad Real. Spain

**IHE AP. Paris, June 2<sup>nd</sup>, 2012**

# Background

SNOMED CT useful in:

- Reference terminology: Information Systems (IS) integration (semantic interoperability)
  - EHR – Pathology IS
    - Electronic request Pathology Study (specimens, anatomy)
    - Pathology report available in EHR (diagnosis, clinical problems, episode coding-billing)
  - Pathology IS and telepathology portal (specimens and diagnosis)
  - Pathology IS and DICOM image manager (specimens and diagnosis)
  - Pathology IS and hospital tumor registry, biobank IS,...
- Interphase terminology
  - Searches made easier, Concept navigation, Qualifiers can be added, synonyms are available

# Pathology reports

- Free text reports
  - Coding specimens (procedures?), anatomy and diagnosis
- Structured reports
  - Coding both the question and the answer
    - Question: Observable entity (Histological grade?)
    - Answer: Diagnosis and qualifiers

CATÁLOGO DE MUESTRAS (SEAP)

C	T	E	SNOMED CT	AUTOPSIAS (A)	Unidades	Minutos	Horas	
1	1	1	56417000	Autopsia adulto, completa	84	840	14	procedure
1	2	1	41770000	Autopsia adulto (sin SNC)	72	720	12	
1	3	1	90864005	Autopsia de alto riesgo	102	1020	17	
1	4	1	74348008	Autopsia parcial, regional, sólo SNC	42	420	7	
1	5	1	309502007	Autopsia fetal (Desde 12 semanas hasta 22 semanas de gestación y/o 500 g)	42	420	7	specimen
1	6	1	4447001	Autopsia perinatal (Desde 22 semanas de gestación y/o 500 g hasta 28 días)	72	720	12	procedure
1	7	1	16361008	Autopsia postnatal (Desde 28 días de vida hasta 1 año de vida)	72	720	12	
1	8	1	430339001	Autopsia pediátrica (Desde 1 año de vida hasta 18 años)	84	840	14	

C	T	E	SNOMED CT	BIOPSIAS (B)	Unidades	Minutos	
2	1	1	309066003	DER Piel, enfermedades dermatológicas, biopsia	6	60	specimen
2	1	2	309075001	DER Piel, lesiones benignas no complejas, exéresis	2	20	
2	1	3	309504008	DER Piel, tumores (punch, afeitado), biopsia	4	40	
2	1	4	35646002	DER Piel, tumores, resección	6	60	procedure
2	2	1	309209004	DIG Esófago, biopsia endoscópica	4	40	specimen
2	2	2	254573006	DIG Esófago, polipectomía	4	40	clinical finding specimen
2	2	3	122630008	DIG Esófago, esofaguectomía	6	60	
2	2	4	438351003	DIG Esófago, esofagogastrectomía	12	120	

# Coding specimens

- Spanish Society of Anatomic Pathology (SEAP): Consensus from 8 institutions from all over Spain.
- Criteria for classifying specimens (without consideration of workload or diagnosis)
  1. General topography (skin)
  2. Obtaining procedure or intention: In biopsies
    - Superficial biopsy (punch, tru-cut or needle)
    - Incisional biopsy
    - Enucleation
    - Complete excision (generally with adjacent tissue)
    - Partial resection of an organ
    - Complete removal of an organ
    - Removal of organ and adjacent structures (e.g. lymphadenectomy)
    - Special resection (Whipple, abdominoperineal,...) o method (stereotactic, endoscopy)

# Coding specimens

- Cytology
  - Exfoliative cytology, brushings or scrapings
  - Fluid sample
  - Touch preparation
  - Fine Needle Aspiration (FNA)
- Autopsy
  - Coded as procedures

# Results of coding specimens and procedures with SNOMED CT

- Biopsies: 345 specimens and procedures
  - 112 specimens (32%) post-coordinated
  - 47 procedures (14%) post-coordinated
- Cytologies: 206 specimens and procedures
  - 73 specimens (35%) post-coordinated
  - 92 procedures (46%) post-coordinated
- Autopsies: 21 different procedures (3 post-coordinated)

# Guidelines for coding with SNOMED CT

- What to do when 2 similar codes are found (choose the most specific one).
  - 128157004 |tissue specimen from brain (specimen)| better than  
119398007 |specimen from brain (specimen)|
  - 438351003 | tissue specimen obtained by esophagogastrrectomy (specimen) | includes:  
122631007 | specimen from esophagus obtained by esophagogastrrectomy (specimen) |
- “esophageal **biopsy sample**” or “**tissue specimen** from esophagus”? “tissue specimen” was preferred



# Guidelines for coding specimens with SNOMED CT

- **specimen from \_\_ (topography) \_\_ obtained by \_\_ (procedure) \_\_**  
*Where procedure <> “biopsy”. Procedure can be general (“excision”) or very specific (“Whipple resection | subtotal adrenalectomy”). (229 concepts)*
- **specimen from \_\_ (topography) \_\_ obtained by \_\_ ( |core |excisional |image guided core |incisional |open |...) \_\_ biopsy**  
*Very specific procedures, from “incisional biopsy” to “stereotactically guided core needle biopsy” or “CT guided needle biopsy”. (55 concepts).*
- **specimen from \_\_ (topography) \_\_ obtained by biopsy (specimen)**  
*Biopsy type is not stated. (46 concepts)*
- **(topography) \_\_ biopsy sample**  
*(75 concepts).*
- **tissue specimen from \_\_ (topography) \_\_**  
*(62 concepts)*

# Postcoordination rules

Refinement (**C:A=V**) (**C:A<sub>1</sub>=V<sub>1</sub>,A<sub>2</sub>=(V<sub>2</sub>),...,A<sub>n</sub>=V<sub>n</sub>**)

- Specimens:

- **specimen : specimen procedure (attribute) = procedure**
- specimen : specimen source topography (attribute) = body structure
- specimen :  
specimen source topography (attribute) = body structure ,  
specimen procedure (attribute) = procedure

- Procedures:

- procedure : procedure site - Direct (attribute) = body structure

# Very specific anatomic regions

- Very few precoordinated terms are very specific (post-coordinated needed)
  - 122683000 | specimen from left kidney, inferior pole obtained by partial nephrectomy (specimen) |
- A specific topographic code may be found as precoordinated:
  - 12212007 | structure of superior segment of left lower lobe of lung (body structure) |

# Laterality

**118169006 | specimen source topography (attribute) |**  
**272741003 | laterality (attribute) |**

Values admitted by this attribute are descendants from the concept: 182353008 | **side (qualifier value) |**:

- 7771000 | left |
  - 51440002 | right and left |
  - 419161000 | unilateral left |
- 24028007 | right |
  - 51440002 | right and left |
  - 419465000 | unilateral right |

# Implementation of specific sites and laterality

- It may be easier implementing a database field “topography” coded with SNOMED CT than post-coordinating specimens or procedures with this attribute.

# A possible error in SNOMED CT

- “122622007 | specimen from pleura obtained by lymph node biopsy |”
- (included as a descendant of “pleura biopsy sample”)

# Adding new concepts

MS Access file

(BatchRequestGeneral\_MSSSI\_Spain\_06032012.mdb)

- Submit Date
- Submitter ID
- Proposed FSN
- Parent ConceptID
- Parent FSN
- Submitter Comment
- Editor Action
- "Reject": Reason
- "Added": UUID
- "Added with Name Change": FSN
- Name Change Reason
- "Found": ConceptID
- "Found": FSN
- Editor Comment

# Coding Pathology diagnosis

- Pathologists should become aware of the structure and contents of SNOMED CT
- A shift from T, P, M schema to a polihierarchy and relationship schema
- Pathologist' diagnosis should be coded as morphology (abnormal body structure) or as clinical diagnosis?



# CAP Anatomic Pathology Subset

<u>Hierarchy</u>	<u>Approximate Count</u>
Body structure	3,805
Clinical Finding	2,024
Procedure	780
Observable entity	386
Specimen	264
Pharmaceutical / Biologic product	83
Record artifact	67
Organism	61
Staging and scales	33
Substance	24
Qualifier value	21
Special concept	12
Events	9
Situation with explicit context	9
Physical object	4
<b>TOTAL CONCEPTS</b> (Jan 2010)	<b>7,582</b>

**In daily practice, this subset was not considered useful by pathologists, mainly due to the mixture of different hierarchies (clinical findings, body structure, observations,...), and due to the fact that many basic pathology diagnosis were not included in the CAP AP Subset.**

# Implementation of SNOMED CT in Hospital General de Ciudad Real

- IHE: Structured Reports Value Sets (1,840 possible values for observations)
- HGUCR: 2,320 pathology diagnosis codes mapped to SNOMED CT
- Search for descriptions that match the local legacy codes was performed using CliniClue Xplorer version 2010 1.243, using SNOMED CT 2011-01-31 International Release and 2010-04-30 Spanish Edition.

# Results. Implementation of SNOMED CT. Hierarchies of precoordinated terms

- Postcoordination needed in 19% (3 conceptID needed only rarely) (81% precoordinated)
- Morphology – Abnormal body structure
  - Only 44% of the codes! (1076)
- Clinical findings
  - 48% of the codes
- Qualifiers, normal anatomic structures, procedures, physical object, substance, organism
  - 10% of the codes

# Results

- 48% of the total terms could be coded using the disorder hierarchy, using:
  - In 1104 terms, using “disorder” hierarchy, e.g. Rosacea
  - In 71 terms, using “finding” hierarchy, e.g. World Health Organization (WHO) grade I (central nervous system tumor)

# Results

- There were a 6% of the local morphological diagnosis that were not well represented using either morphological abnormality or disorder/finding hierarchies, and the following SNOMED CT were found useful (generally combined with other hierarchies):
  - In 95 local terms, the use of a SNOMED CT qualifiers (generally combined with clinical or morphological codes) was found useful, e.g. Granulomatous
  - In 68 local terms, the use of SNOMED CT body structures was found useful, e.g. Undescended testis
  - ...

# Postcoordination in practice

- Initially, in case a combination of terms was needed, the initial combination of terms was made without considering SNOMED CT rules for post-coordination, but considering the best combination of terms that define the concept according to pathologists' idea.

# Attributes

<p><b>Body Structure</b></p> <p><b>Laterality</b></p> <p><b>Clinical Finding</b></p> <p>Associated With</p> <ul style="list-style-type: none"><li>• After</li><li>• Causative Agent</li><li>• Due To</li></ul> <p>Associated</p> <p><b>Morphology</b></p> <p>Clinical Course</p> <p>Episodicity</p> <p>Finding Informer</p> <p>Finding Method</p> <p>Finding Site</p> <p>Has Definitional</p> <p>Manifestation</p> <p>Has Interpretation</p> <p>Interprets</p> <p>Occurrence</p> <p>Pathological Process</p> <p>Severity</p>	<p><b>Events</b></p> <p>Associated With</p> <ul style="list-style-type: none"><li>• After</li><li>• Causative Agent</li><li>• Due To</li></ul> <p>Occurrence</p> <p><b>Evaluation Procedure</b></p> <p>Component</p> <p>Has Specimen</p> <p>Measurement Method</p> <p>Property</p> <p>Scale Type</p> <p>Time Aspect</p> <p><b>Pharmaceutical/Biological Product</b></p> <p>Has Active Ingredient</p> <p>Has Dose Form</p> <p><b>Physical Object</b></p> <p>Has Active Ingredient</p>	<p><b>Procedure</b></p> <p>Access</p> <p>Direct Substance</p> <p>Has Focus</p> <p>Has Intent</p> <p>Method</p> <p>Priority</p> <p>Procedure Device</p> <ul style="list-style-type: none"><li>• Direct Device</li><li>• Indirect Device</li><li>• Using Access Device</li><li>• Using Device</li></ul> <p>Procedure</p> <p>Morphology</p> <ul style="list-style-type: none"><li>• Direct</li><li>• Indirect</li></ul> <p>Morphology</p> <p>Procedure Site</p> <ul style="list-style-type: none"><li>• Procedure Site – Direct</li><li>• Procedure Site – Indirect</li></ul> <p>Recipient Category</p>	<p>Revision Status</p> <p>Route of Administration</p> <p>Surgical Approach</p> <p>Using Energy</p> <p>Using Substance</p> <p><b>Situation with Explicit Context</b></p> <p>Associated Finding</p> <p>Associated Procedure</p> <p>Finding Context</p> <p>Procedure Context</p> <p>Subject Relationship</p> <p>Context</p> <p>Temporal Context</p> <p><b>Specimen</b></p> <p><b>Specimen Procedure</b></p> <p><b>Specimen Source Identity</b></p> <p><b>Specimen Source Morphology</b></p> <p><b>Specimen Source Topography</b></p> <p><b>Specimen Substance</b></p>
--	--	--	--

# Postcoordination

- Morphological concepts (**morphologically abnormal structure**) are **primitive terms** (not “sufficiently” defined) and CANNOT be postcoordinated
- CILLIATED CELL ADENOCARCINOMA
- 35917007 + 125550007
  - 35917007 | Adenocarcinoma, no subtype (morphologic abnormality)
  - 125550007 | Tubal metaplasia (morphologic abnormality)



# An example of lack of coherence in SNOMED CT

- The concept “well differentiated adenocarcinoma” is not available as pre-coordinated term in SNOMED CT in either body structure or in clinical finding hierarchies. However, there are available some more specific similar codes like “follicular adenocarcinoma, well differentiated (morphologic abnormality)” or “well differentiated adenocarcinoma, gastric foveolar type (morphologic abnormality)”.

# Spanish Subset for Diagnosis

- Specimen and Procedure subset is published in <http://www.seap.es/>
- The Spanish Society of Anatomic Pathology is currently working in a pathology diagnosis subset that will be published next October.

# Pathology IS and SNOMED CT

[SNOMED 2	Description SNOMED 2	[Concept	]SNOMED 2
M02140	VALVULA PULMONAR	61699006	M-02140
M02160	VIRUS (HERPES)	36394000	M-02160
M02180	LEUCEMA MIELOIDE AGUDA MINIMAMENTE DIFERENCIADA	112628005	M-02180
M02200	LINFOMA DE CELULAS NK BLASTICAS (9727/3)	62984006	M-02200
M02220	TRICODISCOMA	63068002	M-02220
M02240		71706002	M-02240
M02260		9113006	F-44160
M02300		11498006	M-02300
M02500		65476007	M-02510
M02510		15454001	M-02520
M02520		19776001	M-02530
M02530		4128009	M-02540
M02530		19167000	F-74100
M02530		59540003	F-A8850
M02540		4128009	M-02540
M02550		44832008	M-02521
M02550		1703007	F-44182
M02550		32819004	F-44180
M02550		24122006	M-02541
M02570		65124004	M-02570
M02580		73238002	M-02532
M02590		78558002	M-02590
M02600		14028006	M-02610
M02610		89977008	M-02620
M02610		85345005	F-44300
M02610		111982005	F-44410
M02620		29143009	M-02630
M02620		17719008	F-44302
M02620		63829008	F-44412
M02630		111983000	F-44414
M02700		86991001	M-02710

The screenshot displays the SNOMED CT Navigator interface. On the left, a search bar contains the term 'ACANTOMA'. Below it, a list of search results is shown, including 'acantoma de células claras (trastorno)'. The right pane provides a detailed view of the selected concept, showing its FSN (acantoma de células claras (trastorno)) and preferred name (acantoma de células claras). A table of relationships is also visible, listing various relationships such as 'es un(a)', 'sitio del hallazgo', 'morfología asociada', and 'episodicidad'.



Hoja de Trabajo

En seguimiento

Recientes

+ Hoy (0)

+ Ayer (5)

+ Martes (1)

+ Lunes (0)

+ La semana pasada (7)

+ Hace dos semanas (4)

Procesos

Proced. Diagnóstico

Abreviaturas de Diagnósticos

Técnicas

Fijadores

Estudios

Definición de Protocolos

Definición de Informes

Laboratorios

Catálogo de Procedimientos

Grupos de Trabajo

SNOMED

Estado de Estudios

Estado Bloques

Notas Bloques

Notas Adicionales

Categorías de seguimiento

Parametrización

Regímenes

Pacientes

Personal

Procedencia

Almacén

Modificar técnicas

Bienvenido, MARCIAL GARCÍA ROJO

Hoy es Jueves, 16 de junio de 2011

PAT-Win v.4.0.3 - Build 006

SNOMED 2

Mapeo SNOMED 2 - SNOMED CT

Subconjuntos SNCT

Extensiones

Miembros de extensiones

Actualización SNCT

Nivelación SNCT

# Structured pathology reports

- The general approach to coding items on the CAP and IHE checklist that ask questions has been to use concepts from the SNOMED CT *Observable Entity* hierarchy.
- The answers to these questions can be chosen from concepts in any of several hierarchies. Primarily, they are found in the *Finding*, *Disorder*, or *Morphologic Abnormality* hierarchies (van Berkum, 2003).

# Why we like morphological codes

- As pathologists, we feel the need to have our pathology reports encoded using not only observables (questions) and qualifiers (value sets), but mainly using morphological codes that summarize all observations described in the report.
- Therefore, the section of the pathology dedicated to coding should contain a list of morphological codes associated with the corresponding specimen.

# Clinical findings preferred by EHR

- In **electronic health record** studies, the **clinical findings** should generally be preferred over the morphologic abnormality concept. In case there is only a morphologic abnormality concept, it should be post-coordinated with a focus concept of "64572001|Disease (disorder)|" and linked with the concept model attribute "116676008|Associated morphology (attribute)|" (Lee, 2010).



# Precoordinated primitive terms

- Whenever a pre-coordinated concept is available, that should be the first choice. If the pre-coordinated concept is a primitive concept, an alternative is to create a post-coordinated expression to ensure the necessary semantics are recorded. The reason is that terms encoded with primitive concepts are more difficult to query and test for equivalency (Lee, 2010).



# Linkage concepts in Pathology

- The concept “histological grade (attribute)” is a linkage concept.
- There are some special concepts, like attributes (e.g. 246229000 | histological grade (attribute)), that can be used as “linkage concepts”:
- Question: 371469007 | histologic grade (observable entity),  
246229000 | histological grade (attribute)
- Answer (values):
  - 370114008 | histological grades (qualifier value):
    - 384812005 | moderately differentiated
    - 263843001 | poorly differentiated
    - 263918006 | undifferentiated
    - 263933003 | well differentiated

# Linkage concepts

- Associated morphology is one of the linkage concepts (attributes) used to define clinical findings concepts. Associated morphology attributes specify the morphologic changes seen at the tissue or cellular level that are characteristic features of a disease.

# Morphology attributes

- SNOMED CT concept model provides constraints for attributes that are used as defining relationships, both in distributed SNOMED CT content (so-called pre-coordinated definitions) and in post-coordinated expressions. This model describes that the **domain** of the attribute Associated Morphology is the **Clinical Finding hierarchy**.
- **Procedure hierarchy** concepts can have a Procedure Morphology attribute.
- **Specimen concepts** can have a Specimen Source Morphology attribute.
- All other hierarchies (body structure, events, physical objects,..) cannot have morphology related attributes.

# Morphology attributes

- Permissible values for Associated Morphology, Procedure Morphology, and Specimen Source Morphology attributes are codes belonging to the Morphologically abnormal structure (SCTID 49755003) and its descendants in the Body structure hierarchy.

# Pathological process linkage concept

- When the underlying pathological process for a disorder are not structural and cannot be represented by the associated morphology attribute (e.g. Autoimmune), the pathological process linkage concept can be used. Since July 2009, two new values Infectious process (qualifier value) and Parasitic process (qualifier value) have been added to the range for pathological process.

# **SNOMED CT and digital images: DICOM**

- Specimen Obtained (types, anatomic location, collection procedure)
- Specimen processing (sampling procedure, preparation procedure, stains, fixatives, embedding)
- Diagnosis...

# Open questions

- Should pathology diagnosis be coded using the morphological abnormality sub-hierarchy (part of the Body Structure hierarchy) or should they be coded using Clinical Finding hierarchy?
- Should we try to use mainly morphological codes, and when they are not available, could some diagnoses in pathology reports be coded using clinical (disorder or finding hierarchies)?
- Does using a mixture of hierarchies (Body Structure and Clinical Finding) in pathology diagnosis coding make sense when implementing data exploitation of information systems?
- Should we ask IHTSDO to complete the list of morphological codes to add all those codes missing in the morphological abnormality sub-hierarchy?

# Licensing SNOMED CT

The use of SNOMED CT in production systems requires a [license](#).

- IHTSDO: Most IHTSDO Member countries provides SNOMED CT free of charge to users, although some may charge a small fee for cost recovery. The specific information is provided by your [National Release Centre](#).
- **Qualifying Research Project.** If you are in a country that is not yet a Member of IHTSDO, you can obtain a license to use SNOMED CT through IHTSDO. Qualifying research projects may qualify for a free license.
- Healthcare providers and others wanting to use SNOMED CT in the [countries categorized as low income economies](#) by the World Bank can obtain SNOMED CT Affiliate Licenses free of charge.



# Conclusions

- SNOMED CT needs to be improved in morphology hierarchy
- Pathologists need to understand better the SNOMED CT works. SEAP has published a guideline for specimens and procedures
- Pathology IS allow both local terms and mapping to SNOMED CT
- Digital slides management can be improved with SNOMED CT